



NRW e.V.
Verband allein erziehender
Mütter und Väter



Kinderarmut verringern – Beistandschaft stärken

Gar keiner oder zu wenig Unterhalt sind für viele Alleinerziehende ein erster Schritt in die Armut.

Zahlt der getrennt lebende Elternteil keinen Unterhalt, hilft das Jugendamt im Fachdienst Beistandschaft. Alleinerziehende können sich dort umfassend beraten und unterstützen lassen, um den Anspruch des Kindes durchzusetzen. Jeder kann sich Beistand holen, wenn sie oder er sie braucht! Wie die Beistandschaft arbeitet, haben wir in einem Video erklärt!

Die 5 besten Tipps zur Beistandschaft

the 1990s, the number of people with diabetes has increased in all industrialized countries. In the Netherlands, the prevalence of diabetes is estimated to be 6.5% in 1995, which corresponds to 1.5 million people (1).

Diabetes is a chronic disease, and the long-term consequences of the disease are determined by the degree of glycaemic control. The most serious complications of diabetes are cardiovascular disease, nephropathy, retinopathy, and neuropathy. The prevalence of these complications is directly related to the duration of the disease and the degree of glycaemic control (2).

The aim of this study was to evaluate the effect of a patient education programme on the glycaemic control of patients with diabetes. The programme was designed to improve the patient's knowledge and understanding of the disease, and to encourage the patient to take an active role in the management of the disease.

The programme was evaluated in a randomized controlled trial. The results of the trial are presented in this paper.

Methods

Study design

The study was a randomized controlled trial. The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.

The patients were recruited from a general practice in the Netherlands. The patients were randomized to either the intervention group or the control group. The intervention group received a patient education programme, and the control group received no such programme.



MEHR ALS 1 MIO. KINDER

VON ALLEINERZIEHENDEN

OHNE UNTERHALT.

**HOLEN SIE
SICH
BEISTAND!**

Bildnachweis: a_sro / photocase.de



NRW e.V.
Verband allein erziehender
Mütter und Väter

Das Unterhaltsrecht ist kompliziert. Wir haben fünf Tipps zusammengestellt, auf die Sie bei der Beantragung der Beistandschaft unbedingt achten müssen.

[weiter](#)

Umfrage Beistandschaft und Unterhalt

Gemeinsam mit dem VAMV Bundesverband führten wir 2014 eine Online-Umfrage unter Alleinerziehenden zu den Themen Beistandschaft und Unterhalt durch. Mehr als 1.200 Alleinerziehende nahmen daran teil. Die Auswertung ergab zum Teil alarmierende Ergebnisse: Nur ein Drittel aller Kinder erhielten verlässlichen (Mindest)-Unterhalt.

[weiter](#)